arXiv:1702.06849v1 [math.RT] 22 Feb 2017

Piecewise Hereditary Incidence Algebras

Eduardo do N. Marcos[a], Marcelo Moreira

[a]*Dto de Matemática, Instituto de Matemática e Estatística, Universidade São Paulo, Rua do Matão 1010, Cidade Universitária, CEP 05508-090, São Paulo-SP, Brasil*

Abstract

We present a study of incidence algebras that are piecewise hereditary, which we denominate Phia algebras.

For the Phia algebras $K\Delta$ which are of sheaf type, we study the canonical sincere $K\Delta$-module M and the one-point extension algebra $K\Delta[M]$. We show that if KQ/I is a sincere, quasi-tilted of sheaf type, representation-infinite algebra, then the sincere KQ/I-modules are exceptional. This conclusion allows the construction of a wide range of Phia algebras of wild type of the form $K\Delta[M]$. We explore the simply conectedeness of Phia algebras, we give a positive answer of the so called Skowroński problem for $K\Delta$ a Phia algebra of type $\mathcal{H}$, with connected quiver of tilting objects $\mathcal{K}_{D^b(\mathcal{H})}$: the group $H^1(K\Delta)$ is trivial if, and only if, $K\Delta$ is a simply connected algebra. We determine an upper bound for the strong global dimension of Phia algebras; furthermore, we extend this result for sincere algebras proving that the strong global dimension of a sincere piecewise hereditary algebra is less or equal to three.

Keywords: incidence algebra, piecewise hereditary algebra, simply connected, strong global dimension.
2010 MSC: 13D05, 13D09, 14F05, 16D10, 16E10, 16E35, 16G20

1. Introduction

Throughout the paper K denotes a fixed algebraically closed field. By an algebra we will mean a finite dimensional basic associative K-algebra, a module is always a finite dimensional right module.

Incidence algebras were introduced in the mid-1960s as a natural way of studying some combinatorial problems. In this work, we focus the study on incidence algebras $K\Delta$ associated with a finite poset Δ over K.

The purpose of this paper is to study of incidence algebras that are piecewise hereditary, which we call Phia algebras, *piecewise hereditary incidence algebras*. In addition, given algebras A and B, we use the notation $A \cong' B$ to say that A is derived equivalently to B, that is, $\mathcal{D}^b(A) \cong \mathcal{D}^b(B)$, as triangulated categories.

Email addresses: enmarcos@ime.usp.br (Eduardo do N. Marcos), celovg@ime.usp.br (Marcelo Moreira)

The paper is organized as follows: Section 2 is devoted to fixing the notation and briefly recalling the necessary concepts and results about incidence algebras and piecewise hereditary algebras. In section 3, we explore the simply conectedeness of Phia algebras in order to prove so called Skowroński problem:

Is A simply connected if and only if $H^1(A) = 0$?

We give a positive answer for Phia algebras $K\Delta \cong' \mathcal{H}$, if the quiver of tilting objects of $\mathcal{K}_{D^b(\mathcal{H})}$ is connected:

Theorem. *Let $K\Delta$ be a Phia algebra such that $K\Delta \cong' \mathcal{H}$, whose oriented quiver $\mathcal{K}_{D^b(\mathcal{H})}$ of tilting objects is connected. The group $H^1(K\Delta)$ is trivial if, and only if, $K\Delta$ is simply connected.*

We note that the theorem holds for Phia algebras of quiver type, that is when $K\Delta \cong' KQ$.

The section 4 is dedicated to the study of the global dimension of Phia algebras. In particular, we show that the representation-finite Phia algebras have global dimension less than or equal to two.

Skowroński, Happel and Zacharia have introduced a new homological property called the strong global dimension, in section 5, we determine a upper bound of the strong global dimension of Phia algebras. Moreover, we extend this result to sincere algebras with the theorem:

Theorem. *Let A be a sincere and piecewise hereditary algebra. Then, A have a strong global dimension less or equal to three.*

Section 6, is dedicated to the study of the Phia algebras $K\Delta \cong' \operatorname{Coh}\mathbb{X}$. In this context, we study the $K\Delta$-canonical sincere module M and the one-point extension algebra $K\Delta[M]$. We show that if KQ/I is a sincere, quasi-tilted of quiver sheaf type, representation-infinite algebra, then the KQ/I sincere canonical module is exceptional. This conclusion allows the construction of a wide range of Phia algebras wild type $K\Delta[M]$.

2. Preliminaries

In this section we will recall some definitions which we need. For more details in the subject of incidence algebras we refer to [**?**], [7], [2], for the subject of piecewise hereditary algebras we refer to [4], [27], [6], [1], [8], [9], [30], [12], [13], [14], [15].

There are several equivalent ways of defining incidence algebras, next describe one way for incidence algebras of finite posets.

Definition 2.1 (incidence algebra). Let $(\Delta, \preceq)$ be a poset with n elements. The incidence algebra $K\Delta$ can be viewed as a quotient of the path algebra of the following quiver Q. The vertex set Q_0 is formed by the vertices associated with the points of poset Δ and Q_1 by the arrows α of a vertice a to a vertice b, whenever $a \preceq b$ and there is no $a \preceq c \preceq b$, with $c \neq a$ and $c \neq b$.

Let I be the ideal generated by all commutativity relations $\gamma - \gamma'$, with γ and γ' parallel paths. The incidence algebra $K\Delta$ is KQ/I.

Next, we begin a quick review of the definition of derived category from an abelian category $\mathcal{A}$. First, we recall that the category $\mathcal{K}(\mathcal{A})$ is the quotient of the category of the complexes of $\mathcal{A}$ with its ideal formed by the morphisms homotopic to zero.

The *derived category* $\mathcal{D}(\mathcal{A})$ is obtained from $\mathcal{K}(\mathcal{A})$ via localization with respect to the set of quasi-isomorphisms, that is, from the morphisms $f\colon X \to Y$ of the category of the complexes on $\mathcal{A}$, in which the morphisms induced by the cohomologies $H^i(f)\colon H^i(X) \to H^i(Y)$ are isomorphisms, for each integer i.

Throughout this paper, when we are referring to a bounded derived category of the category $\operatorname{mod} A$ of a K-algebra A, we denote by $\mathcal{D}^b(A)$. Moreover, we assume that the K-category $\mathcal{A}$ is abelian, skeletically small, connected, and Ext-finite (all the spaces of morphisms and the spaces of extensions $\operatorname{Ext}^i(M,N)$ have finite dimensions over K, for every M and N objects of $\mathcal{A}$).

An abelian categories category $\mathcal{H}$ is called *hereditary* if the extensions groups $\operatorname{Ext}^n_{\mathcal{H}}(X,Y) = 0$ for $n \geq 2$ for any pair of objects X and Y of $\mathcal{H}$.

Definition 2.2 (piecewise hereditary category). An abelian category $\mathcal{A}$ is piecewise hereditary of type $\mathcal{H}$ if $\mathcal{H}$ is a hereditary abelian category with a tilting object such that $\mathcal{D}^b(\mathcal{A})$ and $\mathcal{D}^b(\mathcal{H})$ are equivalent as a triangulated categories.

When we say that an algebra A is piecewise hereditary, we mean that the category $\operatorname{mod} A$ is piecewise hereditary. Moreover, there exist a tilting object T in $\mathcal{H}$ such that $A = (\operatorname{End}_{\mathcal{H}} T)^{op}$. The definition of tilting object is as follows:

Definition 2.3 (tilting object [16]). Let $\mathcal{H}$ be a hereditary abelian Ext-finite K-category. The object $T \in \mathcal{H}$ is called tilting if

i) $\operatorname{Ext}^1_{\mathcal{H}}(T,T) = 0$, and
ii) for every $X \in \mathcal{H}$ the condition $\operatorname{Ext}^1_{\mathcal{H}}(T,X) = 0 = Hom_{\mathcal{H}}(T,X)$ implies that $X = 0$.

The above definition was inspired by tilting module definition, which is the following.

Definition 2.4 (tilting module). Let A be a finite dimension algebra over K. An A-module T is called a tilting module if satisfies the following conditions:

i) $\operatorname{Ext}^1(T,T) = 0$,
ii) $\operatorname{pd} T \leq 1$, and
iii) there exists a short exact sequence $0 \to A \to T^1 \to T^2 \to 0$ with $T^1, T^2 \in \operatorname{add} T$.

We state one of the most important theorems about piecewise hereditary algebras.

Theorem 2.5 (Happel [6])**.** *Let $\mathcal{H}$ be an abelian hereditary connected K-category with tilting object. Then $\mathcal{H}$ is derived equivalent to* $\operatorname{mod} KQ$ *or derived equivalent to* $\operatorname{Coh}\mathbb{X}$ *for some weighted projective line $\mathbb{X}$.*

Consequently, we call the algebras $A \cong' KQ$ of piecewise hereditary algebras of quiver type or of type Q. Also, the algebras $A \cong' \operatorname{Coh}\mathbb{X}$ we call piecewise hereditary algebras of sheaf type.

3. Simply connected

In 1993, in the article [17], Skowroński proposed the following problem:

> Given a class of algebras A it is true that A is simply connected if and only if $H^1(A) = 0$, for A in this class?

We conjecture that the statement above is valid for Phia algebras:

Conjecture. *Let $K\Delta$ be a Phia algebra, then $H^1(K\Delta) = 0$ if and only if $K\Delta$ is simply connected.*

The implication $K\Delta$ is simply connected then $H^1(K\Delta) = 0$ has already been proved for incidence algebras in general, De la Peña and Saorín showed the following result:

Theorem 3.1 (De la Peña, Saorín [18])**.** *Let $K\Delta$ be an incidence algebra and (Q, I_v) a presentation of $K\Delta$. Then*

$$\mathrm{Hom}(\pi_1(Q, I_v), K^+) \cong H^1(K\Delta).$$

On the other hand, in the implication "if $H^1(K\Delta) = 0$ then $K\Delta$ is simply connected" it is important to add the hypothesis piecewise hereditary in the incidence algebra, that is, $K\Delta$ to be a Phia algebra. We can see in the example of the projective plane, whose triangulation has the simplicial complex with the fundamental group isomorphic to $\mathbb{Z}_2$ for a field characteristic distinct from 2. Knowing that the fundamental group of the simplicial complex is isomorphic to the fundamental group of poset Δ associated with this complex, applying the previous theorem:

$$\mathrm{Hom}(\mathbb{Z}_2, K^+) \cong H^1(K\Delta).$$

Hence $K\Delta$ is not simply connected but $H^1(K\Delta) = 0$.

We will show that for a large class of Phia algebras the statement "if $H^1(K\Delta) = 0$ then $K\Delta$ is simply connected", is valid.

The main inspiration was the article "Topological invariants of piecewise hereditary algebras" [14] from Le Meur. From the results of Le Meur, we will need the construction of a Galois covering and its relation with the derived equivalences. In order to show this we use the fact that the oriented graph of tilting objects (defined below) is connected.

Definition 3.2 (tilting graph [19]). The oriented graph $\mathcal{K}_{\mathcal{H}}$ of tilting objects of a category $\mathcal{H}$ has the its vertices set in bijection with the isoclasses of the tilting objects. Let T,T' be a non-isomorphisms tilting objects, there exist an arrow $T' \to T$ if $T' = U \oplus Y$, $T = U \oplus X$ with X,Y are non-isomorphisms indecomposables and there is a short exact sequence

$$0 \longrightarrow Y \longrightarrow \tilde{U} \longrightarrow X \longrightarrow 0$$

with $\tilde{U} \in \mathrm{add}\, U$.

In the theorem 3.4, we observe the existence of the hypothesis that the graph $\mathcal{K}_{D^b(\mathcal{H})}$ is connected. Almost all the hereditary categories $\mathcal{H}$ have the connected tilting graph $\mathcal{K}_{D^b(\mathcal{H})}$ ([19], [20], [21]). The case of the category $\mathrm{Coh}(\mathbb{X})$ of wild type is an open problem [21]. The result is valid for all other cases.

Before stating the theorem 3.4, we need the following result:

Lemma 3.3 (Le Meur [14]). *Let $A \cong' \mathcal{H}$ such that $\mathcal{K}_{D^b(\mathcal{H})}$ is connected. Fix a Galois covering $F\colon \mathcal{C} \to A$ with group G and $\mathcal{C}$ locally bounded. Let $T \in D^b(A)$ be a tilting object, $B = \mathrm{End}_{D^b(A)} T$ where $T = T_1 \oplus \ldots \oplus T_n$ is a decomposition in indecomposable objects.*

Let $\lambda_i\colon F_\lambda \widetilde{T}_i \to T_i$ be an isomorphism where $\widetilde{T}_i \in D^b(\mathrm{mod}\,\mathcal{C})$ is indecomposable for every i. This defines the bounded complex of $\mathcal{C}$-modules $\widetilde{T} := \bigoplus_{i,g} {}^g\widetilde{T}_i$ where $g \in G$ and $i \in \{1, \ldots, n\}$.

Let $\mathcal{C}'$ be the full subcategory o $D^b(\mathrm{mod}\,\mathcal{C})$ with objects the complexes ${}^g\widetilde{T}_i$ (for $g \in G$ and $i \in \{1, \ldots, n\}$).

Then the triangle functor $F_\lambda\colon D^b(\mathrm{mod}\,\mathcal{C}) \to D^b(A)$ induces the following connected Galois covering with group G:

$$\begin{aligned} F_{\widetilde{T},\lambda}\colon \mathcal{C}' &\longrightarrow B \\ {}^g\widetilde{T}_i &\longmapsto T_i \\ {}^g\widetilde{T}_i \xrightarrow{u} {}^h\widetilde{T}_j \longmapsto T_i &\xrightarrow{\lambda_j F_\lambda(u) \lambda_i^{-1}} T_j, \end{aligned}$$

The strategy of the proof of the theorem 3.4 is the construction of a Galois covering of $B = \mathrm{End}_{D^b(K\Delta)} T$ from a Galois covering of $D^b(K\Delta)$, where T is a tilting object. This construction will start from a tilting object of $D^b(K\Delta)$. We will show that this tilting object of $D^b(K\Delta)$ has special properties that we will call galoisian. Moreover, the connectivity of the tilting graph of $D^b(\mathcal{H})$ will be important to demonstrate that all tilting objects of $D^b(K\Delta)$ have these special properties.

Let $F\colon \mathcal{C} \to K\Delta$ with group G and $\mathcal{C}$ locally bounded. The G-action in $\mathrm{mod}\,\mathcal{C}$ naturally defines a G-action in $D^b(\mathrm{mod}\,\mathcal{C})$. The push-down functor F_λ induces an exact functor $F_\lambda\colon D^b(\mathrm{mod}\,\mathcal{C}) \to D^b(K\Delta)$ by Le Meur [14], Proposition 2.1. We say that a tilting object has the *galoisians property* the following conditions hold:

- (H_1) for every indecomposable direct summand X of T, there exists $\widetilde{X} \in D^b(\mathrm{mod}\,\mathcal{C})$ such that $F_\lambda \widetilde{X} \cong X$ in $D^b(K\Delta)$;
- (H_2) $\widetilde{X} \ncong {}^g\widetilde{X}$ for every indecomposable direct summand X of T and $g \in G\backslash\{1\}$;
- (H_3) Let $\psi\colon K\Delta \to K\Delta$ an automorphism (and therefore a Galois covering with trivial group), then $\psi_\lambda\colon \mathrm{mod}\,K\Delta \to \mathrm{mod}\,K\Delta$ is an equivalence. So it induces a triangle equivalence $\psi_\lambda\colon D^b(K\Delta) \to D^b(K\Delta)$. If $\psi(x) = x$ for all object x, in $mod(K\Delta)$ then $\psi_\lambda X \cong X$ in $D^b(K\Delta)$ for every indecomposable direct summand X of T.

Theorem 3.4. *Let $K\Delta$ be a Phia algebra such that $K\Delta \cong' \mathcal{H}$. If $\mathcal{K}_{D^b(\mathcal{H})}$ is connected and $H^1(K\Delta) = 0$, then the algebra $K\Delta$ is simply connected.*

Proof. The first step is to consider any Galois covering $F\colon \mathcal{C} \to K\Delta$ a Galois covering with group G, and $\mathcal{C}$ of locally bounded category of finite global dimension.

Suppose $K\Delta \cong' \mathcal{H}$ and $\mathcal{K}_{D^b(\mathcal{H})}$ is connected, by Le Meur [14], Proposition 4.1, Proposition 4.6 and Proposition 4.7, we have that any tilting object $D^b(K\Delta)$ has the galoisian properties. The tilting object satisfying the galoisian properties is essential to use the lemma 3.3 above.

Next, Le Meur constructs a Galois covering to $B = \mathrm{End}_{D^b(K\Delta)} T$ with the same group G of the Galois covering of $K\Delta$ in lemma 3.3 above.

Now we can conclude our theorem by dividing in two cases depending on the type of hereditary abelian category $\mathcal{H}$. Recall:

1. A triangular algebra is simply connected if and only if it has no proper Galois covering.
2. If $K\Delta$ is an incidence algebra, then all its presentations yield isomorphic fundamental groups.

Suppose $\mathcal{H} = \mathrm{mod}\, KQ$ where Q is a finite connected quiver without oriented cycles. Since $K\Delta \cong' \mathcal{H}$, there exists a tilting object T in $D^b(K\Delta)$ such that $\mathcal{H} = \mathrm{End}_{D^b(K\Delta)} T$. By lemma 3.3, the algebra KQ admits a Galois covering with isomorphic group to the fundamental group of $K\Delta$. Since $H^1(K\Delta) = H^1(KQ) = 0$, we get that KQ is tree type, therefore simply connected. Then the fundamental group of $K\Delta$ is trivial.

The other case is when $\mathcal{H}$ has no nonzero projective object. Hence there exist a squid algebra S such that $\mathcal{H} \cong' S$. Consequently, there exist a tilting object T in $D^b(K\Delta)$ such that $S = \mathrm{End}_{D^b(K\Delta)} T$. By lemma 3.3, the algebra S admits a Galois covering with isomorphic group to the fundamental group of $K\Delta$. By the article of Le Meur [14], $H^1(K\Delta) = H^1(S) = 0$, then S is simply connected. We conclude that the fundamental group of $K\Delta$ is trivial. □

4. Global dimension

The *projective dimension* of an object M in an abelian category $\mathcal{A}$ is by definition

$$\mathrm{pd}_{\mathcal{A}} M = \sup\{d \in \mathbb{N} \,/\, \mathrm{Ext}^d_{\mathcal{A}}(M, M') \neq 0 \text{ for some } M'\}.$$

and the *global dimension* of $\mathcal{A}$ is the $\sup\{\mathrm{pd}_{\mathcal{A}} M : M \text{ is an object in } \mathcal{A}\}$.

When we refer to the global dimension of algebras A, we are considering the global dimension of the category $\mathrm{mod}\, A$. The piecewise hereditary algebras have finite global dimension. We mention the following result:

Theorem 4.1 (Happel-Reiten-Smalø [3])**.** *Let $\mathcal{A}$ be a piecewise hereditary, abelian category with finite length and n non-isomorphisms simples objects. Then the category $\mathcal{A}$ has the global dimension less than or equal to n.*

An interesting question is whether there is a special upper bound for Phia algebras? We found a positive answer in the article "Bounds on the global dimension of certain piecewise hereditary categories" [7] of Ladkani. He obtained another upper bound for the global dimension of piecewise hereditary algebras. Before we state Ladkani's theorem, some concepts are needed. Let $G(\mathcal{A})$ be the directed graph whose vertices are the isomorphism classes of indecomposables of $\mathcal{A}$, and there exist a arrow $M \longrightarrow M'$ between the vertices M and M', if $\mathrm{Hom}_{\mathcal{A}}(M, M') \neq 0$. In this graph, a walk between two vertices is a sequence $\epsilon = (\epsilon_0, \ldots, \epsilon_{r-1})$, in $\{-1, 1\}^r$ and r is an integer greater than or equal to one. An ϵ-path from M to M' is a sequence of vertices $M_0 = M$, M_1,...,$M_r = M'$ such that

$$M_i \longrightarrow M_{i+1} \text{ in } G(\mathcal{A}) \text{ if } \epsilon_i = 1, \text{and}$$
$$M_{i+1} \longrightarrow M_i \text{ in } G(\mathcal{A}) \text{ if } \epsilon_i = -1.$$

Theorem 4.2 (Ladkani [7])**.** *Let $\mathcal{A}$ be a finite length, piecewise hereditary category. Assume that there exist $r \geq 1$, $\epsilon \in \{-1, 1\}^r$ and an indecomposable M_0 such that, for any indecomposable M, there exists an ϵ-path from M_0 to M. Then*

$$\mathrm{gldim}\,\mathcal{A} \leq r + 1 \quad \textit{and} \quad \mathrm{pd}_{\mathcal{A}}\, M + \mathrm{id}_{\mathcal{A}}\, M \leq r + 2, \quad \textit{for any indecomposable } M.$$

We note that the hypothesis piecewise hereditary is necessary, for example, the result is not valid for the class of local algebras which, in general, have an infinite global dimension. One of the consequences of the theorem concerns the so-called *sincere algebra*. We recall that an algebra is called sincere if it admits a sincere indecomposable module in its category of modules, that is, an indecomposable module such that all simple modules occur as factors of their composition series. Incidence algebra are sincere as we see next.

Proposition 4.3. *Let $K\Delta = KQ_{K\Delta}/I$ be an incidence algebra. Then $K\Delta$ is a sincere algebra.*

Proof. We need to show the existence of a module M indecomposable sincere about $K\Delta$. The candidate M is the following module representation:

a) for each vertice a in $Q_{K\Delta}$ we associate K;
b) for each arrow $\alpha\colon a \to b$ in $Q_{K\Delta}$ We associate the identity $1\colon K \to K$.

First, we will show that M is indecomposable. For this, we will study the $\mathrm{End}\, M$. We consider $f = (f_a)_{a \in Q_0}$ a non-zero morphism of $\mathrm{End}\, M$. Thus, there exists $f_a\colon K \to K$ not zero for some $a \in Q_0$, implying that f_a is an isomorphism. Given an arrow $\alpha\colon a \to b$, we have that $f_a 1 = 1 f_b$. If the arrow is in the other direction, we get the same result. Thus, no matter the direction of the arrow, we conclude that $f_a = f_b$. So, since the graph is connected, we always have a walk connecting the vertex a to any vertex c:

By a finite process, we conclude that $f_a = f_c$ for every vertex c of $Q_{K\Delta}$ and consequently $\operatorname{End} M \cong K$. Then M is an indecomposable module.

It is clear that M is sincere.

Therefore, M is an indecomposable sincere module of the incidence algebra $K\Delta$. □

Definition 4.4. We call the module M in the proposition above the *canonical sincere module.*

We confirm that Phia algebras are sincere, so the following result applies to them.

Corollary 4.5 (Ladkani [7])**.** *Let A be a piecewise hereditary, sincere, finite-dimensional algebra. Then* $\operatorname{gldim} A \leq 3$ *and* $\operatorname{pd} M + \operatorname{id} M \leq 4$*, for any indecomposable module M in* $\operatorname{mod} A$*.*

The representation-finite sincere algebras have global dimension less than or equal to two. Before proving this affirmation, we need the definition of a directed module.

Definition 4.6 (directed module). Consider a cycle in the module category, that is a sequence:

$$M_0 \xrightarrow{f_1} M_1 \xrightarrow{f_2} M_2 \xrightarrow{f_3} \ldots \to M_{t-1} \xrightarrow{f_t} M_t \cong M_0\,,$$

where f_i are non-zero morphisms between indecomposables and non-isomorphic modules, for $1 \leq i \leq t$.

Let M be an um indecomposable module, M is called directed if it does not belong to any cycle.

Happel showed in the article "On the derived category of a finite-dimensional algebra" the following result:

Corollary 4.7 (Happel [22])**.** *Let A be a representation-finite, piecewise hereditary algebra. Then* $\operatorname{mod} A$ *is directed, that is, all A-modules are directed.*

For a directed algebra A, given an indecomposable M in $\operatorname{mod} A$, if there is a non-isomorphism, non-zero $M \xrightarrow{f} M$ then M belongs to a cycle. This contradicts the fact that A is directed. Thus, we observe that $\operatorname{End} M$ is isomorphic to K for any indecomposable A-module M.

Now, we use the following Ringel theorem [23]:

Theorem 4.8 (Ringel [23])**.** *Let A be an algebra having a sincere and directed indecomposable module. Then A is a tilted algebra.*

We prove the next result:

Proposition 4.9. *Let A a representation-finite sincere algebra. Then A is a tilted algebra, consequently* $\operatorname{gldim} A \leq 2$*.*

Proof. We need to show that $\operatorname{mod} A$ has a sincere and directed indecomposable module. We know $\operatorname{mod} A$ has a canonical sincere module. Since A is a representation-finite sincere algebra, this sincere indecomposable A-module is directed, thanks to Happel's result 4.7.

We come to the conclusion that A is a tilted algebra, therefore it has global dimension dimension less than or equal to two. □

Another connection we can make with the global dimension is with the *strongly simply connected incidence algebras*. Before we show the results visualizing this connection, we will introduce a family of algebras with a global dimension equal to three called critical algebras.

Definition 4.10 (critical algebra [24]). Let B be an algebra. We say that B is critical if it satisfies the following properties:

(i) B has a unique source i and a unique sink j,

(ii) Let S_a be a simple module associated with source a and let S_b be a simple module associated with sink b, then $\operatorname{pd} S_a = 3$ and $\operatorname{id} S_b = 3$. If S is a simple module associated with other vertice then $\operatorname{pd} S \leq 2$ and $\operatorname{id} S \leq 2$.

(iii) Consider the minimal projective resolution of the simple module S_a:

$$0 \longrightarrow P_3 \longrightarrow P_2 \longrightarrow P_1 \longrightarrow P_0 \longrightarrow S_a \longrightarrow 0.$$

Give the B-module $P = \bigoplus_{k=0}^{3} P_k$. Then all indecomposable projective are in $\operatorname{add} P$, and each indecomposable projective is a direct summand of exactly one P_k, for $k \in \{0, \ldots, 3\}$.

(iv) Consider the minimal injective resolution of the S_b:

$$0 \longrightarrow S_b \longrightarrow I_0 \longrightarrow I_1 \longrightarrow I_2 \longrightarrow I_3 \longrightarrow 0.$$

Consider the B-module $I = \bigoplus_{k=0}^{3} I_k$. Then all indecomposable injectives are in $\operatorname{add} I$, and each indecomposable injective is a direct summand of exactly one I_k, for $k \in \{0, \ldots, 3\}$.

(v) B does not contain any proper full subcategory that verifies i), ii), iii) e iv).

A description of the quivers with relations of all the critical algebras can be found in the work "A criterion for global dimension two for strongly simply connected schurian algebras" [24].

Proposition 4.11 (Bordino-Fernndez-Trepode [24])**.** *Let B be a critical algebra. Then the algebra has one of the following presentations.*



In case of Q_n, the relations are given by declaring that two parallel paths are equal.

Using this proposition, we see that the only critical incidence algebras are the ones whose quiver is the Q_n, since the presentations of algebras A_1, A_l, B_1 and B_l have non-commutative relations.

Now we can state a theorem of Bordino, Fernández and Trepode [24] which we will use in our proposition 4.14.

Theorem 4.12 (Bordino-Fernández-Trepode [24])**.** *Let A be a strongly simply connected schurian algebra with global dimension greater or equal three. Then there exists a full subcategory B of A such that B is critical.*

Before stating the proposition 4.14, we need the following result:

Theorem 4.13 (Assem-Castonguay-Marcos-Trepode [25]). *Let $K\Delta$ be an incidence algebra, $K\Delta$ is strongly simply connected if and only if $K\Delta$ does not contain a crown.*

Now we have the result on incidence algebras.

Proposition 4.14. *Let $K\Delta$ be a strongly simply connected incidence algebra. Then the global dimension of $K\Delta$ is less than or equal to two.*

Proof. We recall that an incidence algebra is a schurian algebra. If the global dimension of $K\Delta$ is greater than or equal to three then it will contain a full subcategory B that is critical. Since $K\Delta$ is an incidence algebra, B is of the form Q_n and thus, the incidence algebra $K\Delta$ would contain a crown. This contradicts the result above.

The result is proved □

5. Strong global dimension

Let $\mathcal{P}_A$ be a full subcategory of $\operatorname{mod} A$ consisting of the projective modules. We denote $C^b(\mathcal{P}_A)$ the category of bounded complexes with entries in $\mathcal{P}_A$ and $K^b(\mathcal{P}_A)$ the corresponding homotopic category. If $P = (P^i, d^i) \in K^b(\mathcal{P}_A)$ is not zero, there are $r \leq s$ such that $P^i = 0$ for all $i < r$ and $P^i = 0$ for all $s < i$, with P_r and P_s not zero. The length of P is defined as $\mathrm{l}(P) = s - r$.

Definition 5.1 (strong global dimension [26]). Let A a finite-dimensional algebra. Then, we define the strong global dimension of A by

$$\operatorname{sgldim} A = \sup\{\mathrm{l}(P) \mid P \in K^b(\mathcal{P}_A) \text{ indecomposable}\}.$$

Observe that the strong global dimension can be infinite.

The next theorem has a importance to the study of the strong global dimension for Phia algebras.

Theorem 5.2 (Happel-Zacharia [26]). *Let A a finite-dimensional algebra. The algebra is piecewise hereditary if and only if* $\operatorname{sgldim} A$ *is finite.*

Thus it is clear that Phia algebras have finite strong global dimension. A tool to compute the strong global dimension of an algebra is the following theorem.

Theorem 5.3 (Alvares, Le Meur, Marcos [15]). *Let $\mathcal{T}$ be triangulated category which is triangle equivalent to the bounded derived category of a hereditary abelian category. Let $T \in \mathcal{T}$ a tilting object.*

There exists a full and additive subcategory $\mathcal{H} \subset \mathcal{T}$ which is hereditary and abelian, such that the embedding $\mathcal{H} \hookrightarrow \mathcal{T}$ extends to a triangle equivalence $D^b(\mathcal{H}) \cong \mathcal{T}$, and

$$T \in \bigsqcup_{m=0}^{l} \mathcal{H}[m]$$

for some integer $l \geq 0$. Moreover, if $\operatorname{End}_{\mathcal{T}} T)^{op}$ is not hereditary then there exists such a pair $(\mathcal{H}, l)$ verifying $\operatorname{sgldim}(\operatorname{End}_{\mathcal{T}} T)^{op} = l + 2$

Inspired by article "Jordan Hölder theorems for derived module categories of piecewise hereditary algebras" [12], Alvares, Le Meur and Marcos have used an alternative definition of the strong global dimension [15].

Lemma 5.4 (Alvares, Le Meur, Marcos [15])**.** *Let $T \in D^b(\mathcal{H})$ be a tilting object such that $A = (\operatorname{End} T)^{op}$. Given a object Y in $D^b(\mathcal{H})$, we define*

$$\ell_T^+(Y) = \sup\{n \in \mathbb{Z} \mid \operatorname{Hom}_{D^b(\mathcal{H})}(Y, T[n]) \neq 0\}$$
$$\ell_T^-(Y) = \inf\{n \in \mathbb{Z} \mid \operatorname{Hom}_{D^b(\mathcal{H})}(T[n], Y) \neq 0\}.$$

Then the strong global dimension of A is $\operatorname{sgldim} A = \sup\{\ell_T^+(Y) - \ell_T^-(Y) \mid Y \in D^b(\mathcal{H})\}$.

We want to give an upper bound for the strong global dimension of the Phia algebras. We have a more general result.

Theorem 5.5. *Let A be a sincere, piecewise hereditary algebra. Then*

$$\operatorname{sgldim} A \leq 3 .$$

Proof. We consider M the sincere module and $A = P_1 \oplus \ldots \oplus P_n$ the decomposition of A in indecomposable modules.

By hypothesis, there exist a quasi-inverse functor $F\colon D^b(A) \to D^b(\mathcal{H})$ which makes the triangulated equivalence, where $\mathcal{H}$ is a hereditary category.

Using the sincere module property, for each $i \in \{1, \ldots, n\}$ it follow:

$$0 \neq \operatorname{Hom}_A(P_i, M) = \operatorname{Hom}_{D^b(A)}(P_i, M) \cong \operatorname{Hom}_{D^b(\mathcal{H})}(FP_i, FM).$$

Let $FA = T$ be a tilting object in $D^b(\mathcal{H})$ such that $(\operatorname{End} T)^{op} \cong A$. For each $i \in \{1, \ldots, n\}$, we denote $FP_i = T_i[m_i]$ the indecomposable direct summand of T where $T_i \in \mathcal{H}[0]$ such that $D^b(\mathcal{H}) = \bigsqcup_{m \in \mathbb{Z}} \mathcal{H}[m]$. Also we denote $FM = M'[m]$ where $M' \in \mathcal{H}[0]$.

Now, for each $i \in \{1, \ldots, n\}$, we have the following:

$$0 \neq \operatorname{Hom}_{D^b(\mathcal{H})}(FP_i, FM) = \operatorname{Hom}_{D^b(\mathcal{H})}(T_i[m_i], M'[m]).$$

Therefore $m_i = m$ or $m_i = m - 1$, for each $i \in \{1, \ldots, n\}$, implying that the indecomposables direct summands of T are in $\mathcal{H}[m-1]$ or $\mathcal{H}[m]$ and shows that $\operatorname{sgldim} A \leq 3$. □

As an immediate consequence of previous theorem, we get the following:

Corollary 5.6. *Let $K\Delta$ be a Phia algebra . Then* $\operatorname{sgldim} K\Delta \leq 3$.

Another corollary is what has already been proved by Ladkani in [7].

Corollary 5.7. *Let A be a sincere, piecewise hereditary algebra. Then*

$$\operatorname{gldim} A \leq 3 .$$

Proof. This is a consequence of inequality $\operatorname{gldim} A \leq \operatorname{sgldim} A$. □

Example 5.8. *The incidence algebra, whose quiver is given below, is a Phia algebra which has global dimension equal to three, and therefore it also has strong global dimension three.*



By the result 4.14, the family of Phia algebras that has the global dimensional and strong global dimension equal to three are not strongly simply connected. Can this family be more characterized?

6. Phias of canonical type

In 1987, Geigle and Lenzing introduced the category $\operatorname{Coh}\mathbb{X}$ called category of coherent sheaves on the weighted projective line $\mathbb{X}$ [27]. This category has an important relation with the category of modules over a canonical algebra, $\operatorname{Coh}\mathbb{X}$ is derived equivalent to category of modules over a canonical algebra $C(p,\lambda)$ [27]. Some references to this type of algebra are: "Tame algebras and integral quadratic forms" [23] and "Elements of the representation theory of associative algebras, volume three" [28].

The article "A class of weighted projective curves arising in representation theory of finite dimensional algebras" [27] by Geigle and Lenzing, and the paper "Introduction to coherent sheaves on weighted projective line" [4] by Chen and Krause are important texts for an introduction to the theory of category of coherent sheaves on weighted projective line. We will use the characterization of the category $\operatorname{Coh}\mathbb{X}$ exposed in "Hereditary noetherian categories with a tilting complex" [8] by Lenzing.

The Phia algebras $K\Delta \cong' \operatorname{Coh}\mathbb{X}$, were studied in the article "Which canonical algebras are derived equivalent to incidence algebras of posets?" [1] of Ladkani, which in part, relates to our work. This article was published in 2008, and presents a direct study of Phia algebras of sheaves type or Phia algebras of canonical type, nomenclature influenced by the derived equivalence between the categories $\operatorname{Coh}(\mathbb{X})$ and the category of modules over a canonical algebra $C(p,\lambda)$. The main theorem of this article follows:

Theorem 6.1 (Ladkani [1])**.** *Let $K\Delta$ be an incidence algebra. If $C(p,\lambda) \cong' K\Delta$ then $p = (p_1,p_2,p_3)$ or $p = (p_1,p_2)$, where $p_i \geq 2$. For each weighted projective line with $p = (p_1,p_2,p_3)$ or $p = (p_1,p_2)$, where $p_i \geq 2$, there exist a incidence algebra $K\Delta$ such that $C(p,\lambda) \cong' K\Delta$.*

We shall use the notation $\mathrm{Coh}(\mathbb{X}) = \mathcal{C}$.

The previous categories $\mathcal{C}$ are divided into 3 disjoint classes that are: quiver, tubular and wild. These classes are characterized according to the Euler characteristic $\mathcal{X}(\mathbb{X})$ of the weighted projective lines, let $\mathbb{X} = (p_1, \ldots, p_n; \lambda_3, \ldots, \lambda_n)$, the Euler characteristic is defined as follows:

$$\mathcal{X}(\mathbb{X}) = 2 - \sum_{i=1}^{n}(1 - \frac{1}{p_i}).$$

If $\mathcal{C}$ has the Euler characteristic greater than zero then $\mathcal{C}$ is of quiver type, that is, it is derived equivalent to category of modules over a hereditary algebra of extended Dynkin type. If $\mathcal{C}$ has the Euler characteristic equal to zero then it is of the tubular family, the wild family has the negative Euler characteristic.

For the purpose of exibiting some families of Phia algebras of sheaves type, we will study one-point extension. The works of Barot, De la Peña and Lenzing ([10], [11] and [29]) provide conditions for piecewise hereditary algebras A and for the A-modules M that result in piecewise hereditary algebras $A[M]$.

Example 6.2. *Let M be the sincere canonical module 4.3 and consider the one-point extension $K\Delta[M]$.*



We observe that $K\Delta[M]$ is an incidence algebra that has a poset with a unique maximal element represented by the vertex $*$. The one-point extension by the canonical sincere module M refers us to a technique used in the study of posets, to add a unique maximal element.

We denote $\mathcal{C}_+$ (resp. $\mathcal{C}_0$) the full subcategory of all vector bundles (resp. sheaves of finite length) on $\mathbb{X}$.

Assume that we have a triangular equivalence:

$$F\colon D^b(K\Delta) \to D^b(\mathcal{C}).$$

We were inspired by the work "Quasi-tilted algebras of canonical type" [9] and show that the canonical sincere module M is associated, by F, with a bundle. Lenzing and Skowroński characterize each indecomposable module over a representation-infinite, quasi-tilted algebra of sheaf type.

Now we can state a theorem of Lenzing and Skowroński [9] which we will use in our proposition 6.4.

Proposition 6.3 (Lenzing-Skowroński [9]). *Let B be a quasi-tilted of sheaf type, representation-infinite algebra. Then each indecomposable B-module belongs to one of the subcategories*

a) $\mathrm{mod}_0^l(B)$ *consisting of all* $Y[-1]$ *where* Y *in* $\mathcal{C}_0'$ *satisfies* $\mathrm{Hom}_{\mathcal{C}}(T_+, Y) = 0$ *and* $\mathrm{Ext}^1_{\mathcal{C}}(T_0', Y) = 0$*;*

b) $\mathrm{mod}_+(B)$ *consisting of all* Y *from* $T_0''^{\perp} \cap \mathcal{C}_+$ *satisfying* $\mathrm{Ext}^1_{\mathcal{C}}(T_+, Y) = 0$*;*

c) $\mathrm{mod}_0^c(B)$ *consisting of all* Y *from* $\mathcal{C}_0$ *satisfying* $\mathrm{Hom}_{\mathcal{C}}(T_0', Y) = 0$ *and* $\mathrm{Ext}^1_{\mathcal{C}}(T_0'', Y) = 0$*;*

d) $\mathrm{mod}_-(B)$ *consisting of all* $Z[1]$ *with* Z *in* $T_0'^{\perp} \cap \mathcal{C}_+$ *satisfying* $\mathrm{Hom}_{\mathcal{C}}(T_+, Z) = 0$ *;*

e) $\mathrm{mod}_0^r(B)$ *consisting of all* $Z[1]$ *with* Z *in* $\mathcal{C}_0$ *satisfying* $\mathrm{Hom}_{\mathcal{C}}(T_+, Y) = 0$ *and* $\mathrm{Hom}_{\mathcal{C}}(T_0'', Y) = 0$.

Proposition 6.4. *Let $A = KQ/I$ be a sincere, quasi-tilted of sheaf type, representation-infinite algebra. Assume that M is an indecomposable sincere module and $F\colon D^b(A) \to D^b(\mathcal{C})$ a quasi-inverse functor which makes the triangulated equivalence. Then FM is a bundle.*

Proof. By hypothesis, $D^b(A)$ is equivalent as a triangulated category to $D^b(\mathcal{C})$, so there is a tilting object T in $D^b(\mathcal{C})$ that $(\mathrm{End}_{D^b(\mathcal{C})}(T))^{op} = A$.

Since M is sincere, we have $\mathrm{Hom}_A(P_x, M) \neq 0$ for each vertice x of Q. Thus, given a vertice x of Q, using the functor F we obtain

$$\mathrm{Hom}_A(P_x, M) \cong \mathrm{Hom}_{D^b(A)}(P_x, M) \cong \mathrm{Hom}_{D^b(\mathcal{C})}(FP_x, FM).$$

Let $FP_x = P_x'[p]$ and $FM = Y[m]$, where P_x', Y belongs to $\mathcal{C}[0]$. Therefore,

$$\mathrm{Hom}_{D^b(\mathcal{C})}(FP_x, FM) \cong \mathrm{Hom}_{D^b(\mathcal{C})}(P_x'[p], Y[m]).$$

Hence, $\mathrm{Hom}_{D^b(\mathcal{C})}(P_x'[p], Y[m])$ is not zero.

$P_x'[p]$ is an indecomposables direct summands of T, where T decompose into a direct sum $T_0'[-1] \oplus T_+ \oplus T_0''$, see [9]. Let G be the quasi-inverse of F

$$G\colon D^b(\mathcal{C}) \to D^b(A) \quad \text{ such that } GT = A,$$

we get that $FP_{x_1} \oplus \ldots \oplus FP_{x_n}$ is isomorphic to $T = T_0'[-1] \oplus T_+ \oplus T_0''$, for each $x_i \in Q$ where $i = 1, \ldots n$. That is, $P_{x_i}'[p]$ is isomorphic to an indecomposable direct summand of T consequently $p = 0$ or $p = -1$.

We use the proposition of Lenzing-Skowroński [9] above. Take $x \in Q$ such that $FP_x = P_x'[0]$ and $\mathrm{Hom}_{D^b(\mathcal{C})}(P_x', FM) \neq 0$ then FM is not isomorphic to $Y[-1]$ eliminating the case a). Analogously, by the group of morphisms $\mathrm{Hom}_{D^b(\mathcal{C})}(P_x'[-1], FM) \neq 0$ then FM is not isomorphic to $Y[1]$ eliminating the cases d) and e). The cases b) and c) such that $FM \cong Y$. We assume that the case c) is valid to the module M. Then $\mathrm{Hom}_{\mathcal{C}}(T_0', FM) = 0$, that is, $\mathrm{Hom}_{D^b(\mathcal{C})}(P_x', FM) = 0$ for some $x \in Q$ contradicting the fact that M is sincere. Therefore, the case c) is not valid for M. Hence $FM \in T_0''^{\perp} \cap \mathcal{C}_+$, then FM is bundle. □

We remember that Phia algebras are sincere. The next step is to demonstrate that the sincere canonical module M is exceptional. We recall the definition of exceptional object:

Definition 6.5 (exceptional). An object E in a triangulated K-category $\mathcal{T}$ is called exceptional if $\mathrm{End}(E) = K$ and, E not have auto-extensions, that is, $\mathrm{Hom}_{\mathcal{T}}(E, E[n]) = 0$ for each non-null integer n.

Corresponding, let A a finite dimensional K-algebra, the A-module E is called exceptional if $\mathrm{End}(E) = K$, and $\mathrm{Ext}^1_A(E, E) = 0$.

Again, the next proposition is not specific to incidence algebras.

Proposition 6.6. *Let $A = KQ/I$ be a sincere, quasi-tilted algebra of quiver-sheaf type and representation-infinite. Assume M an indecomposable sincere module. Then M is exceptional.*

Proof. We have a triangulated equivalence:

$$F\colon D^b(A) \to D^b(\mathcal{C}).$$

We show that $\mathrm{Ext}^1_{\mathcal{C}}(FM, FM) = 0$, we have that

$$\mathrm{Ext}^1_{\mathcal{C}}(FM, FM) = \mathrm{Hom}_{D^b(\mathcal{C})}(FM, FM[1]) \cong \mathrm{Ext}^1_A(M, M).$$

By proposition 6.4, FM in $\mathcal{C}_+$. Therefore, FM is a bundle. Then we use the theorem of Lenzing-Reiten [30] which states that if $\mathcal{C}$ of quiver type, then each indecomposable bundle is exceptional.

Hence, the indecomposable sincere module M is exceptional. □

Before stating the lemma 6.8, we need the following result:

Theorem 6.7 (Lenzing-De la Peña [29])**.** *Let A be derived equivalent to a canonical algebra Λ of weight type $(p_1, \ldots, p_n)$. We fix a triangle-equivalence $D^b(A) \cong D^b(\mathcal{C})$ for the weighted projective line $\mathbb{X}$ corresponding to Λ. Let M be an exceptional A-module.*

Up to translation in $D^b(\mathcal{C})$, the module M corresponds to an exceptional vector bundle over $\mathbb{X}$ and if $\mathcal{X}(\mathbb{X}) > 0$, then $A[M]$ is derived equivalent to the path algebra of a wild connected quiver.

Lemma 6.8. *Let $A = KQ/I$ be a sincere, quasi-tilted of quiver-sheaf type, representation-infinite algebra. Let M be an indecomposable sincere module. Then $A[M]$ is derived equivalent to the path algebra of a wild connected quiver.*

Proof. We have a triangulated equivalence:

$$F\colon D^b(A) \to D^b(\mathcal{C}).$$

By proposition 6.6, M is exceptional. Moreover, by proposition 6.4, M associated, by F, with a exceptional vector bundle over $\mathbb{X}$.

We use the theorem of Lenzing-De la Peña [29] above. We conclude that $A[M]$ is derived equivalent to the path algebra of a wild connected quiver. □

We can produce several examples of Phia algebras where $K\Delta[M]$ is of wild type.

Example 6.9. *Among the Phia algebras of extended Dynkin type, we have chosen the concealed algebras that satisfy all the hypotheses of the lemma 6.8. The example is of a Phia algebra of type* $\widetilde{\mathbb{E}}_6$.

Let $K\Delta$ be a Phia algebra of tubular-sheaf or wild-sheaf type, we are interested in characterize which modules M are such that $K\Delta[M]$ is piecewise hereditary. The works ([10], [11] and [29]) deal with this situations. The most present hypothesis on M is the requirement that M is a derived simple module. We recall the definition.

Definition 6.10 (derived simple). Let X be a indecomposable module over a algebra A of finite global dimension. X is called derived simple, if the following conditions are satisfied:

i) the "middle term" E of the Auslander-Reiten triangle

$$\tau X \to E \to X \to \tau X[1]$$

is indecomposable;

ii) X is τ-periodic in $D^b(A)$.

A natural question to ask now is when the sincere canonical module is exceptional?

Before stating the proposition 6.12 about exceptional sincere canonical module, we need the following result:

Proposition 6.11 (Gastaminza, De la Peña-Platzeck, Redondo, Trepode [31])**.** *Let* $A = KQ/I$ *be a strongly simply connected incidence algebra. Let* M *be the unique indecomposable* A*-module with* $\dim_K M(x) = 1$ *for each* $x \in Q$*. Then*

a) $\operatorname{pd} M \leq 1$*;*
b) $\operatorname{Ext}^1_A(M, M) = 0$.

Proposition 6.12. *Let* $K\Delta$ *a representation-finite Phia algebra of quiver-sheaf or tubular-sheaf type. Let* M *be the canonical indecomposable sincere module. Then is exceptional.*

Proof. The first step is to show that $K\Delta$ is simply connected. This follows from the theorem of Assem-Skowroński, [32], which states that if A is a algebra derived equivalent to tubular canonical algebra or to euclidean hereditary algebra, then A is simply connected.

We get that $K\Delta$ is simply connected.

The second step is to prove that $K\Delta$ is strongly simply connected. This follows from the result of Bretscher-Gabriel [33] which states that a representation-finite algebra is strongly simply connected if, and only if, the algebra is simply connected.

The final step in the proof is to use proposition of Gastaminza-De la Peña-Platzeck-Redondo-Trepode [31] above. Therefore the canonical sincere module M is exceptional. □

We recall that a strongly simply connected Phia algebra is quasi-tilted, by proposition 4.14.

For representation-finite, quasi-tilted Phia algebras $K\Delta$ of quiver-sheaf type, we prove that the canonical sincere $K\Delta$-module M is exceptional and $K\Delta[M]$ is a Phia. Thus, by propositions 6.6 and 6.12, we ensure that M is exceptional only for:

- quasi-tilted $K\Delta$ of quiver-sheaf type;
- quasi-tilted, representation-finite $K\Delta$ of tubular-sheaf type.

Acknowledgements

The first named author has been supported by the thematic project of Fapesp 2014/09310-5.

The second named author acknowledges support from CAPES, in the form of a Phd Scholarship, Phd done at programa de Matemática, IME-USP, (Brazil).

References

References

[1] S. Ladkani, Which canonical algebras are derived equivalent to incidence algebras of posets?, Communications in Algebra 36 (12) (2008) 4599–4606.

[2] J. Bustamante, On the fundamental group of a schurian algebra, Communications in Algebra 30 (11) (2002) 5307–5329.

[3] D. Happel, I. Reiten, S. Smalø, Piecewise hereditary algebras, Archiv der Mathematik 66 (3) (1996) 182–186.

[4] X. Chen, H. Krause, Introduction to coherent sheaves on weighted projective line, techreport, arXiv:0911.4473v3 [math.RT] (Sep. 2010).

[5] D. Happel, J. Rickard, A. Schofield, Piecewise hereditary algebras, Bulletin London Mathematical Society 20 (1) (1988) 23–28.

[6] D. Happel, A characterization of hereditary categories with tilting object, Inventiones mathematicae 144 (2) (2001) 381–398.

[7] S. Ladkani, Bounds on the global dimension of certain piecewise hereditary categories, Journal of Pure and Applied Algebra 212 (9) (2008) 2140–2145.

[8] H. Lenzing, Hereditary noetherian categories with a tilting complex, Proceedings of the American Mathematical Society 125 (7) (1997) 1893–1901.

[9] H. Lenzing, A. Skowroński, Quasi-tilted algebras of canonical type, Colloquium Mathematicum 71 (2) (1996) 161–181.

[10] M. Barot, H. Lenzing, Derived canonical algebras as one-point extensions, Contemporary Mathematics 229 (1998) 7–15.

[11] M. Barot, H. Lenzing, One-point extensions and derived equivalence, Journal of Algebra 264 (1) (2003) 1–5.

[12] L. Hügel, S. Koenig, Q. Liu, Jordan-hölder theorems for derived module categories of piecewise hereditary algebras, Journal of Algebra 352 (1) (2012) 361–381.

[13] D. Zacharia, D. Happel, Homological properties of piecewise hereditary algebras, Journal of Algebra 323 (2009) 1139–1154.

[14] P. Le Meur, Topological invariants of piecewise hereditary algebras, Transactions of the American Mathematical Society 363 (4) (2011) 2143–2170.

[15] E. Marcos, P. Le Meur, E. Alvares, The strong global dimension of piecewise hereditary algebras, arXiv:1305.5213v2 [math.RT].

[16] D. Happel, I. Reiten, Hereditary abelian categories with tilting object over arbitrary base fields, Journal of Algebra 256 (2002) 414–432.

[17] A. Skowroński, Simply connected algebras and hochschild cohomologies, Canadian Mathematical Society: Conference Proceedings 14 (1993) 431–447.

[18] J. De la Peña, M. Saorín, On the first hochschild cohomology group of an algebra, Manuscripta Mathematica (104) (2001) 431–442.

[19] D. Happel, L. Unger, On the set of tilting objects in hereditary categories, Fields Institute Communications 45 (2005) 141–159.

[20] A. Buan, R. Marsh, M. Reineke, I. Reiten, G. Todorov, Tilting theory and cluster combinatorics, Advances in Mathematics 204 (2006) 572–618.

[21] M. Barot, D. Kussin, H. Lenzing, The cluster category of a canonical algebra, Transactions of the American Mathematical Society 362 (8) (2010) 4313–4330.

[22] D. Happel, On the derived category of a finite-dimensional algebra, Commentarii Mathematici Helvetici (62) (1987) 339–389.

[23] C. Ringel, Tame algebras and integral quadratic forms, Vol. 1099 of Lectures Notes in Mathematics, Springer, 1984.

[24] N. Bordino, E. Fernández, S. Trepode, A criterion for global dimension two for strongly simply connected schurian algebras, arXiv:1110.6160v1 [math.RA].

[25] I. Assem, D. Castonguay, E. Marcos, S. Trepode, On strongly simply connected algebras and multiplicative bases, Journal of Algebra 283 (1) (2005) 161–189.

[26] D. Happel, D. Zacharia, A homological characterization of piecewise hereditary algebras, Mathematische Zeitschrift 260 (1) (2008) 177–185.

[27] W. Geigle, H. Lenzing, A class of weighted projective curves arising in representation theory of finite dimensional algebras, Vol. 1273 of Lectures Notes in Mathematics, Springer, 1987.

[28] D. Simson, A. Skowroński, Elements of the representation theory of associative algebras, Vol. 72 of London Math. Soc. Student Text, Cambridge Univ. Press, 2007.

[29] H. Lenzing, J. De la Peña, Spectral analysis of finite dimensional algebras and singularities, Series of Congress Reports, 2008, pp. 541–588.

[30] H. Lenzing, I. Reiten, Hereditary noetherian categories of positive euler characteristic, Mathematische Zeitschrift (254) (2006) 133–171.

[31] S. Gastaminza, J. De la Peña, M. Platzeck, M. Redondo, S. Trepode, Finite dimensional algebras with vanishing hochschild cohomology, Journal of Algebra 212 (1) (1999) 1–16.

[32] I. Assem, A. Skowroński, On some classes of simply connected algebras, Proceedings of the London Mathematical Society s3-56 (3) (1988) 417–450.

[33] O. Bretscher, P. Gabriel, The standard form of a representation-finite algebra, Bulletin de la Société Mathématique de France 111 (1983) 21–40.